

# Properties of complete bipartite codimension two subspace arrangements

Zach Teitler<sup>1</sup>   Douglas A. Torrance<sup>2</sup>

<sup>1</sup>Boise State University

<sup>2</sup>Monmouth College

January 15, 2014

## Linear subspace arrangements

Let  $\mathbb{k}$  be a field. A **linear subspace** of  $\mathbb{P}^n$  is a variety  $V(I)$  where  $I \subset R = \mathbb{k}[x_0, \dots, x_n]$  is an ideal generated by linear forms. If a linear subspace has dimension  $d$ , then we may call it a  **$d$ -plane**. Let  $\mathcal{A}$  be an arrangement of linear subspaces. We define

$$V_{\mathcal{A}} = \bigcup_{X \in \mathcal{A}} X$$

$$I_{\mathcal{A}} = \bigcap_{X \in \mathcal{A}} I(X) = I(V_{\mathcal{A}}).$$

# Minimal graded free resolutions and graded Betti numbers

Given a graded ideal  $I$ , consider a **minimal graded free resolution**

$$0 \rightarrow F_p \xrightarrow{\varphi_p} F_{p-1} \xrightarrow{\varphi_{p-1}} \cdots \rightarrow F_1 \xrightarrow{\varphi_1} F_0 \xrightarrow{\varphi_0} I \rightarrow 0$$

For each  $i$ ,  $F_i$  is a **graded free module**, i.e.,  $F_i \cong \bigoplus_j R(d_j)$

## Definition

The **graded Betti numbers** of  $I$  are

$$\beta_{i,j} = \# \text{ of copies of } R(-j) \text{ in } F_i$$

## Betti tables

We may list all the graded Betti numbers of a minimal graded free resolution using a **Betti table**:

	0	1	...	$i$	...
$\vdots$					
1	$\beta_{0,1}$	$\beta_{1,2}$		$\beta_{i,i+1}$	
2	$\beta_{0,2}$	$\beta_{1,3}$		$\beta_{i,i+2}$	
$\vdots$					
$j$	$\beta_{0,j}$	$\beta_{1,j+1}$		$\beta_{i,i+j}$	
$\vdots$					

# Regularity

## Definition 1

The **(Castelnuovo-Mumford) regularity** of a graded ideal  $I$  is

$$\operatorname{reg} I = \max\{j : \beta_{i,i+j} \neq 0 \text{ for some } i\}$$

Note that this is the index of the last nonzero row of the Betti table.

## Definition 2

If  $I$  is the ideal of a variety, this is equivalent to the following definition using the cohomology of the ideal sheaf.

$$\operatorname{reg} I = \min\{j : h^i(\mathbb{P}^n, \tilde{I}(j-i)) = 0 \forall i > 0\}$$

## Our question

Let  $\mathcal{A}$  be a subspace arrangement. What is the Castelnuovo-Mumford regularity of  $I_{\mathcal{A}}$ ?

## What is known

Theorem (Derksen, Sidman (2002))

If  $\mathcal{A}$  is a linear subspace arrangement, then

$$\operatorname{reg} I_{\mathcal{A}} \leq |\mathcal{A}|$$

This bound is sharp. For example, an arrangement of  $d$  skew lines intersecting a line  $L$  (which is not in the arrangement) in  $d$  distinct points will have a regularity of  $d$ . The Betti table for for the case of 5 such lines in  $\mathbb{P}^3$  is given below.

	0	1
3	1	.
4	6	9
5	9	2

# Incidence graphs

Suppose  $\mathcal{A}$  is a subspace arrangement in  $\mathbb{P}^n$ .

## Definition


The **incidence graph** of  $\mathcal{A}$  is the graph  $\Gamma(\mathcal{A})$  such that

- $V(\Gamma(\mathcal{A})) = \mathcal{A}$
- $E(\Gamma(\mathcal{A})) = \{XY : \dim(X \cap Y) > \text{expdim}(X \cap Y)\}$ .



## Lines in $\mathbb{P}^3$

Two lines in  $\mathbb{P}^3$  can either intersect in a point or not at all.



Note that the example we saw of  $d$  lines with regularity  $d$  above has incidence graph  $dK_1$ , i.e., no edges.

What happens to the regularity when we impose more structure?

# Complete bipartite graphs


## Definition

A graph  $G = (V, E)$  is the **complete bipartite graph**  $K_{a,b}$  if

- $V = V_1 \cup V_2$  where  $|V_1| = a$  and  $|V_2| = b$ .
- If  $u, v \in V_1$  or  $u, v \in V_2$ , then  $uv \notin E$ .
- If  $u \in V_1$  and  $v \in V_2$  or vice versa, then  $uv \in E$ .

## Example

The complete bipartite graph  $K_{2,3}$  is as follows:



## Question

If  $\Gamma(\mathcal{A}) = K_{a,b}$  with  $a \leq b$ , then what is  $\text{reg } I_{\mathcal{A}}$ ?

## Example 1

Using Macaulay 2, we can construct  $(n - 2)$ -plane arrangements with the desired incidence graphs.

If  $n = 3$  and  $\Gamma(\mathcal{A}) = K_{3,3}$ , then  $I_{\mathcal{A}}$  has the following Betti table:

	0	1
2	1	.
3	1	.
4	.	1

$$\text{reg } I_{\mathcal{A}} = 4$$

## Example 2

If  $n = 3$  and  $\Gamma(\mathcal{A}) = K_{5,10}$ , then  $I_{\mathcal{A}}$  has the following Betti table:

	0	1	2
2	1	.	.
3	.	.	.
4	.	.	.
5	.	.	.
6	.	.	.
7	.	.	.
8	.	.	.
9	.	.	.
10	6	10	4


$$\operatorname{reg} I_{\mathcal{A}} = 10$$

## The result

### Theorem

If  $\Gamma(\mathcal{A}) = K_{a,b}$  with  $a \leq b \leq 3$ , or  $3 \leq a \leq b$ , then  $\text{reg } I_{\mathcal{A}} = \max\{a + 1, b\}$ .

*Sketch of proof.* Suppose  $\mathcal{A}$  is a line arrangement in  $\mathbb{P}^3$ . Then  $\mathcal{A}$  consists of rulings of a quadric surface  $Q$ .



## Sketch of proof, cont.

The result follows from computing cohomologies using the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2) \xrightarrow{\cdot Q} \tilde{I}_{\mathcal{A}} \rightarrow \mathcal{I}_{V_{\mathcal{A}} \cap Q, Q} \rightarrow 0$$

For  $n > 3$ , it can be shown that  $V_{\mathcal{A}}$  is a cone over a line arrangement in  $\mathbb{P}^3$  with the same incidence graph, and therefore it has the same regularity. □

# Arithmetic Cohen-Macaulayness


The sheaf cohomology calculations used to prove the above result can also be used to prove the following result.

## Theorem

*If  $\Gamma(\mathcal{A}) = K_{a,b}$  with  $a \leq b \leq 3$ , or  $3 \leq a \leq b$ , then  $V_{\mathcal{A}}$  is arithmetically Cohen-Macaulay if and only if  $b \in \{a, a + 1\}$ .*



Thank you!



*Enrico Belletti*